



Combined ACL and ALL reconstruction reduces the rate of reoperation for graft failure or secondary meniscal lesions in young athletes

Pierre Laboudie¹ · Adil Douiri¹ · Nicolas Bouguennec¹ · Alexandre Biset¹ · Nicolas Gravelleau¹

Received: 30 October 2021 / Accepted: 18 March 2022

© The Author(s) under exclusive licence to European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2022

Abstract

Purpose Graft failure and secondary meniscal tears are major concerns after anterior cruciate ligament (ACL) reconstruction in young athletes. The aim was to evaluate the link between ACL reconstruction with and without anterolateral ligament (ALL) reconstruction and outcomes in young patients participating in pivoting sports.

Methods This was a retrospective study of data collected prospectively. Patients less than 20 years, involved in pivoting sports and undergoing primary ACL reconstruction with a quadruple hamstring tendon (4HT) graft or 4HT graft combined with anterolateral ligament reconstruction (4HT + ALL) were included. Survival analysis was performed to identify the prognostic indicators for reoperation due to graft failure or secondary meniscal lesions. Knee laxity was assessed and patient reported outcome measures (PROMs) were collected.

Results A total of 203 patients (mean (\pm SD) age: 16.3 ± 2 years) with a mean follow-up of 4.8 ± 0.9 (range: 3.3–6.8) years were included. There were 101 4HT and 102 4HT + ALL grafts. Graft rupture rates were 11.9% for 4HT grafts and 5.8% for 4HT + ALL grafts (n.s.). There were 9.9% secondary meniscal procedures for 4HT grafts vs. 1.9% for 4HT + ALL grafts ($p = 0.02$). With reoperation for graft failure or secondary meniscal lesions at final follow-up as the endpoint, survival was better in the 4HT + ALL group (91.4% vs. 77.8%, respectively; $p = 0.03$). Absence of ALL reconstruction (HR = 4.9 [95%CI: 1.4–17.9]; $p = 0.01$) and preoperative side-to-side laxity > 3 mm (HR = 3.1 [95%CI: 1.03–9.1]; $p = 0.04$) were independently associated with an increased rate of reoperations. Mean (\pm SD) side-to-side laxity was 1.3 ± 1.3 mm (range: -2 to 5) for 4HT grafts vs. 0.9 ± 1.3 mm (range: -6 to 4.8) for 4HT + ALL grafts (n.s.) 6 months post-surgery. The rate of return to the same sport at the same level was 42.2% for 4HT grafts vs. 52% for 4HT + ALL grafts (n.s.). There was no significant difference in subjective outcomes including PROMs between the two groups.

Conclusion Combined ALL + ACL reconstruction reduced the rate of graft failure and secondary meniscal injury in young athletes when compared to ACL reconstruction alone. Subjective results were comparable, with a similar rate of complications. Combined reconstruction should be preferred in this young population.

Level of evidence Level IV.

Keywords Knee · Anterior cruciate ligament · Anterolateral ligament · Graft rupture · Return to sport

Introduction

Anterior cruciate ligament (ACL) injuries have become increasingly prevalent in adolescent athletes as sports training has increased in frequency and intensity [17]. Treatment of ACL injuries in this population may be challenging

because of an unacceptably high rate of graft failure and reoperation [14, 34, 43, 53, 57].

Reported rates of graft failure vary from 1.4 to 18% [23, 30, 31, 50, 52], reflecting the fact that the causes of graft failure are multifactorial. This high risk was confirmed by the Norwegian ACL registry [37] showed that age is a significant risk factor for revision with a hazard ratio (HR) = 4.0 for the youngest age group (15–19 years) compared to subjects > 30 years of age. The Swedish ACL registry [2] found that adolescent patients (defined as between 13 and 19 years old) have the highest rates of early revision.

✉ Pierre Laboudie
pierre.laboudie@gmail.com

¹ Clinique du Sport de Bordeaux-Mérignac, 4 rue George Negrevergne Merignac, 33700 Bordeaux, France

It has also been shown that ACL reconstruction may fail to provide full control of laxity and restoration of normal knee kinematics which could therefore increase the risk of secondary meniscal injury [20, 41, 58].

For this reason, there is currently great interest in the role of the anterolateral structures of the knee in controlling laxity and their ability to share loads with the ACL graft and its effect on meniscal healing and return to sport [11, 26, 42, 47]. Ferretti et al. [15] found magnetic resonance imaging (MRI) abnormalities of the ALL in 88.2% of patients with ACL rupture. A systematic review of 29 articles by Ariel de Lima et al. [3] reported that the ALL was found in 100% of cases and that MRI was a good method to visualise the ALL. Mechanoreceptors have also been identified in the ALL, confirming its proprioceptive role [4]. Sonnery-Cottet et al. [50] demonstrated that combined HT + ALL reconstruction was associated with significantly lower rates of graft rupture and secondary meniscal tear [49, 50] in a high-risk population when compared to HT and bone–patellar tendon–bone (B-PT-B) grafts. In addition, Lee et al. [28] observed that adolescents with ACL rupture have greater rotational instability and a lower rate of healing of the ALL than adults.

The aim of this study was to compare the reoperation rates for graft failure and secondary meniscal lesions in patients undergoing a quadruple hamstring tendon (4HT) graft versus a 4HT graft combined with anterolateral ligament reconstruction (4HT + ALL). The secondary aims were to evaluate the prognostic factors for graft failure, defined as reoperation for graft failure or secondary meniscal injury, and to determine the subjective outcomes and patient reported outcome measures (PROMs) between 4 and 4HT + ALL reconstruction. The hypothesis was that combined reconstruction of the ACL and ALL would reduce the rate of graft failure and secondary meniscal injury without causing more complications.

Materials and methods

Study population

This single-centre, single-surgeon, retrospective, cohort study was approved by our institutional review board (CE Clinique du Sport, Mérignac—12-2019-12). Patients undergoing ACL reconstruction with a minimum follow-up of 3 years were identified in our prospective surgical database. A total of 933 knees that underwent primary ACL reconstruction between 2015 and 2018 were reviewed. The inclusion criteria were: primary ACL reconstruction with an isolated 4HT (Group 1) or 4HT + ALL (Group 2); age < 20-years; and athlete practicing a pivoting sport. A pivoting sport was defined as a sport that involves rotational pivoting movements of the knee when the foot is on

the ground (e.g. skiing or tennis); a contact pivoting sport is the same with the possibility of contact between players (e.g. football or rugby). Exclusion criteria included: age > 20-years; absence of practice of a pivoting sport; prior knee surgery; multi-ligament injury; other graft; refusal to participate. The flow of the patients is shown in Fig. 1. There were 240 ACL reconstructions fulfilling the inclusion criteria and 37 patients were lost to follow-up; 203 patients were included in the final analysis: 101 in Group 1 (4HT) and 102 in Group 2 (4HT + ALL).

Surgical technique

All patients underwent intra-articular reconstruction with a 4HT autograft with the semi-tendinosus tendon, according to the single antero-medial bundle biological augmentation (SAMBBA) technique [46], with two suspensory buttons, Pullup and Pullup XL (SBM, Lourdes, France).

ALL reconstruction was then performed with a gracilis graft [11], using a modification of the technique of Boutsidis et al. [7] A femoral tunnel was drilled just posterior and proximal to the lateral epicondyle and another was drilled in the tibia just posterior to the Gerdy tubercle 7 mm below the joint-line (Fig. 2). The ALL graft was then fixed in the femur with a resorbable interference screw (Arthrex, Naples, Florida) and then fixed in the tibia with the ACL tight rope (Arthrex, Naples, Florida), tightened in extension and neutral rotation after ACL tensioning.

The decision to perform ALL reconstruction was based on ALL Expert Group recommendations [45]: pivot shift \geq grade 2, hyperlaxity (knee recurvatum $> 10^\circ$), high-level athlete (competing at regional or national level), Second fracture (bony avulsion of the tibial ALL insertion on radiograph).





Fig. 1 Patient flow through the study. ACL, anterior cruciate ligament; 4HT, quadruple hamstring tendon; 4HT + ALL, quadruple hamstring tendon + anterolateral ligament reconstruction

Fig. 2 **A** Lateral view and **B** anteroposterior view of the 4HT + ALL grafts. The intra-articular short ACL graft is fixed with two suspensory buttons (Pullup and Pullup XL; SBM, Lourdes, France). The ALL graft is fixed into an independent femoral tunnel with a resorbable interference screw (Arthrex, Naples, Florida) and in the tibia with an ACL tight rope (Arthrex, Naples, Florida)



Postoperative protocol

All patients had the same specific rehabilitation protocol to recover mobility, particularly in extension and contraction of the quadriceps [51]. Return to sport was only allowed after 6 months with validation of an isokinetic test, symmetrical jump test, ACL return to sport after injury (RSI) scale > 85% and teaching of a prevention program.

Outcome measures

Radiographic measurement of the tibial slope, measured in relation to the anatomical axis [59], was performed by two knee sport fellows (PL and AD) and showed excellent inter-observer reliability (intra class correlation) = 0.94 [95%CI: 0.92–0.95]. Antero-posterior side-to-side laxity using the GNRB® device [40] (whose threshold was evaluated, respectively, at 3 and 1.5 mm for complete and partial ACL rupture [28, 29]) was performed preoperatively. Intra-operative surgical data (duration of surgery, findings, other procedures, HT graft diameter) were collected.

Patients were assessed at 6 weeks (physical examination), 6 months (physical examination, side-to-side laxity with the GNRB®) and 3 years (physical examination, questionnaire and PROMs). The questionnaire included the following question: (1) about your operated knee, do you experience occasional pain, swelling, stiffness, instability or discomfort due to the surgical hardware? The PROMs collected were the subjective International Knee Documentation Committee

(IKDC), ACL-RSI, Tegner activity scale and Lysholm knee score.

Graft failure was defined as recurrent instability requiring a revision procedure, a recurrent lesion with MRI confirmation of a graft lesion, or side-to-side laxity > 3 mm, while a secondary meniscal lesion was defined as meniscal pain with a confirmed MRI lesion requiring meniscal surgery, whether the meniscal lesion was new or a previous failed meniscal repair. Failure was defined as reoperation for graft failure or secondary meniscal lesions.

Study population

The demographic characteristics of the study population are summarised in Table 1. Mean (\pm SD) age was 16.3 ± 2 years (range: 11–20) and 119 patients (58.8%) were male. The majority of patients ($n = 168$, 82.8%) were involved in a pivot contact sport.

Statistical analysis

Quantitative variables are reported as mean \pm standard deviation (SD) and range (min–max), while qualitative variables are reported as number and percentage.

Nominal variables were compared using the Chi² test or Fisher's exact test depending on the sample size. Variance was tested using the Student's *t* test or Mann–Whitney *U* test when a non-parametric test was required.

Kaplan–Meier survival curves were used to determine survival, with reoperation for graft failure or secondary meniscal tear as an endpoint. A Cox regression analysis was performed

Table 1 Characteristics of the study cohort

	All patients (<i>N</i> =203)	4HT (<i>N</i> =101)	4HT+ALL (<i>N</i> =102)	<i>p</i> -value
Follow-up (years)				< 0.01
Mean ± SD	4.8 ± 0.9	5.4 ± 0.8	4.3 ± 0.7	
Range (min–max)	(3.3–6.8)	(3.5–6.8)	(3.3–6.1)	
Male sex, <i>n</i> (%)	119 (58.8)	57 (56.4)	62 (60.8)	n.s.
Age (years)				n.s.
Mean ± SD	16.3 ± 2	16.5 ± 2.2	16.8 ± 1.9	
Range (min–max)	(11–20)	(11–20)	(12–20)	
BMI (kg/m ²)				n.s.
Mean ± SD	21.9 ± 2.9	21.8 ± 3.0	22.1 ± 3.0	
Range (min–max)	(16–35)	(16–35)	(17–31)	
Time, injury to surgery (months)				n.s.
Mean ± SD	4.9 ± 6.5	4.3 ± 3.8	5.7 ± 8.2	
Range (min–max)	(1–60)	(1–24)	(1–60)	
Sport, <i>n</i> (%)				
Pivot contact	168 (82.8)	78 (77.2)	90 (88.2)	0.04
Pivot non-contact	35 (17.2)	23 (22.8)	12 (11.7)	
Tibial slope (°)				n.s.
Mean ± SD	7.7 ± 2.7	7.7 ± 2.7	7.6 ± 2.8	
Range (min–max)	(1–15)	(1–15)	(1–15)	
Side-to-side pre-surgical laxity (mm)				n.s.
Mean ± SD	3.1 ± 1.9	2.9 ± 1.9	3.4 ± 2.0	
Range (min–max)	(0–9)	(0–9)	(0–8)	
Meniscal tears, <i>n</i> (%)	113 (56)	55 (55)	58 (58.9)	n.s.
Suture, <i>n</i> (%)	93 (45)	48 (47.5)	45 (44.1)	n.s.
Meniscectomy, <i>n</i> (%)	20 (9.9)	7 (6.9)	13 (12.8)	n.s.
Chondral lesions, <i>n</i> (%)	43 (21)	22 (21.8)	21 (20.6)	n.s.
Diameter of the ST graft (mm)				n.s.
Mean ± SD	8.7 ± 0.7	8.6 ± 0.7	8.7 ± 0.6	
Range (min–max)	(7–10)	(7–10)	(7–10)	
Duration of surgery (min)				< 0.01
Mean ± SD	32.8 ± 8.4	28.4 ± 6.9	37.1 ± 7.5	
Range (min–max)	(16–76)	(16–50)	(17–76)	

Significance was set at $p < 0.05$

HT, hamstring tendon; HT + ALL, hamstring tendon + anterolateral ligament; BMI, body mass index; SD, standard deviation; NS, not significant

to the determine factors associated with failure. The variables included in the analysis were selected based on statistically significant results of univariate analysis ($p < 0.05$). Post hoc analysis on the sample size (203 patients) calculation found a power of 84.5% to find a difference in combined graft failure or secondary meniscal injury between the two groups. All statistical analyses were performed using IBM SPSS (Statistical Product and Service Solutions) software for Windows (version 27).

Results

Reoperations

At a final follow-up of 4.8 ± 0.9 years (range: 3.3–6.8), there was a lower rate of graft failure in the 4HT + ALL group compared to the 4HT group, but the difference was not statistically significant (5.8% vs. 11.9%; n.s.). The

rate of secondary meniscal lesions was also lower in the 4HT + ALL group (1.9% vs. 9.9%; $p=0.02$). There were more secondary meniscal procedures in patients who underwent meniscal repair compared to those who did not (7.5% [7/93] vs. 1.8% [2/110] of knees, respectively; $p=0.04$). The details of the reoperations are summarised in Table 2.

Predictors of failure

At the final follow-up, mean (\pm SD) overall survival was $83.3 \pm 3\%$ (Fig. 3). Survival was $77.8 \pm 4.3\%$ in the 4HT group vs. $92.2 \pm 2.7\%$ in the 4HT + ALL group ($p=0.04$) (Fig. 4). The predictors of failure are summarised in Tables 3 and 4.

Table 2 Reoperations in whole cohort and two sub-groups at the final follow-up

Reoperation	All patients (N=203)	4HT (N= 101)	4HT + ALL (N= 102)	p-value
Graft failure	18 (8.9)	12 (11.9)	6 (5.8)	n.s.
Time to graft failure (years)				n.s.
Mean \pm SD	1.9 \pm 1.0	2.0 \pm 1.2	1.8 \pm 0.5	
Range (min–max)	(0.8–4.5)	(0.8–4.5)	(1.2– 2.5)	
Secondary meniscal lesion	12 (5.9)	10 (9.9)	2 (1.9)	0.02
Medial meniscus	9 (4.4)	8 (3.4)	1 (0.5)	0.03
Lateral meniscus	3 (1.5)	2 (1)	1 (0.5)	n.s.
Cyclops	18 (8.9)	10 (9.9)	8 (7.8)	n.s.
Hardware removal	3 (1.5)	0 (0)	3 (2.9)	n.s.
Infection	2 (1)	2 (2)	0 (0)	n.s.

Significance was set at $p < 0.05$

All values shown are n (%) unless stated otherwise

HT, hamstring tendon; HT + ALL, hamstring tendon + anterolateral ligament; SD, standard deviation; NS, not significant ($p > 0.05$)

Fig. 3 Kaplan–Meier analysis of overall graft survival





Fig. 4 Kaplan–Meier analysis of survival based on the surgical procedure



Outcomes

Mean (\pm SD) post-surgical side-to-side laxity was similar in the two groups: 1 ± 1.3 mm in the 4HT + ALL group vs. 1.3 ± 1.3 mm in the 4HT group (n.s.). The answers to the questionnaire and PROMs at 2 years were also similar in the two groups (n.s.) (Table 5).

Discussion

The most important finding of this study is that the combined rate of reoperation for graft failure or secondary meniscal tear was decreased significantly by the addition of ALL reconstruction in young athletes. ALL reconstruction was associated with a 4.9-fold reduction in reoperation for graft failure or secondary meniscal lesions at a mean follow-up of 4.8 years. Recently, several studies have highlighted the role of the anterolateral structures of the knee in the control of rotatory laxity [10, 19, 21, 36, 42]. Song et al. [44] carried out a systematic review of 326 patients in 7 studies evaluating persistent rotatory instability after ACL reconstruction and found a significantly lower prevalence of residual pivot shift in patients treated by lateral extraarticular tenodesis (LET) + ACL reconstruction (13.3%) than in those with ACL reconstruction only (27.2%). It is possible that this better control of rotational laxity protects the knee and consequently reduces the load on both the graft and the menisci (repaired or not). Furthermore, postoperative sagittal side-to-side laxity was similar in the two groups, which is consistent with the findings of Sonnery-Cottet et al. [50];

thus, it could be argued that ALL reconstruction plays a primary role in the control of rotational laxity.

It has not been proved conclusively that ALL reconstruction can decrease graft failure rates. A recent meta-analysis of 6 studies including 683 patients with a mean age of 27 years by Rhatomy et al. [39] found no difference in the rate of graft failure between isolated ACL reconstruction vs. combined ACL + ALL reconstruction. However, the current results are similar to those of Sonnery-Cottet et al. [50] who found a graft failure rate of 4.1% in a group of 16–30-year-olds who had undergone 4HT + ALL, while Lee et al. [27] did not observe any graft failure among 42 ACL revisions with ALL reconstruction. Another recent study by Balendra et al. [5] found an ACL reconstruction failure rate of 8.2% at 2 years in a population of professional footballers with an average age of 23 years, with a decrease in risk of graft failure to 2% when a B-PT-P graft was associated with a LET. In the setting of ACL revision surgery, Rayes et al. [38] also compared HT + ALL to the supposed gold standard B-PT-TB + LET and found the same outcomes regarding graft failure and reoperation, thus demonstrating the strength of this reconstruction.

Concerning the adolescent population studied here, the most interesting comparison is with the meta-analysis of Wiggins et al. [57], who found that the pooled rate of ipsilateral ACL reinjury across 11 studies was 10% in patients <25-years-old, at a mean follow-up of 51 months. Not only is the graft failure rate of the 4HT + ALL group lower, but the age of the patients was also lower. The rest of the literature reports graft failure rates of between 9 and 25% in the same young population with different types of grafts

Table 3 Univariate analysis of predictors of failure at final follow-up

Factor			Mean \pm SD survival (%) at final follow-up (4.8 \pm 0.9 years)	* <i>p</i> -value
Demographics pre-surgical	Age (years)	≤ 15	81.8 \pm 4.8	n.s.
		> 15	83.4 \pm 3.9	
	Sex	Male	80.6 \pm 4.2	n.s.
		Female	87.1 \pm 3.9	
	BMI (kg/m ²)	≤ 25	81.1 \pm 3.4	n.s.
		> 25	93.8 \pm 4.2	
	Tibial slope [12]	≤ 11	82.8 \pm 3.2	n.s.
		> 11	85.7 \pm 7.6	
	Pivot contact sport	Yes	83.6 \pm 3.3	n.s.
		No	82.3 \pm 6.6	
Side-to-side laxity (mm) at 6 months	≤ 3	93.0 \pm 3.0	0.05	
	> 3	80.3 \pm 6.0		
Injury to surgery (months)	≤ 6	82.8 \pm 3.4	n.s.	
	> 6	83.0 \pm 6.7		
Surgical details	Surgical group	HT	77.8 \pm 4.3	0.04
		HT + ALL	92.2 \pm 2.7	
	Meniscal lesion	Yes	77.4 \pm 4.5	0.02
		No	90.7 \pm 3.5	
	Meniscectomy	Yes	73.2 \pm 10.4	n.s.
		No	84.5 \pm 3.0	
	Meniscal repair	Yes	78.6 \pm 4.9	n.s.
		No	87.4 \pm 3.5	
HT graft diameter (mm) [1]	≤ 8	77.2 \pm 6.3	n.s.	
	> 8	85.6 \pm 3.3		

Significance was set at $p < 0.05$

*Log rank test

HT, hamstring tendon; HT + ALL, hamstring tendon + anterolateral ligament; BMI, body mass index; SD, standard deviation; NS, not significant ($p > 0.05$)

Table 4 Multivariate analysis of predictors of failure at final follow-up

Predictor	Hazard ratio \pm SD [95% CI]	<i>p</i> -value
Surgical group—HT	4.9 \pm 0.7 [1.4–17.9]	0.01
Pre-surgical side-to-side laxity > 3 mm	3.1 \pm 0.5 [1.03–9.1]	0.04
Presence of meniscal lesion	2.5 \pm 0.6 [0.8–7.8]	n.s.

Significance was set at $p < 0.05$

HT, hamstring tendon; SD, standard deviation; CI, confidence interval, NS, not significant ($p > 0.05$)

(HT, B-PT-B, Fascia Lata) [6, 18, 24, 52], which are higher than those in the present 4HT + ALL group. The recent stability experience study [16] found that adding a LET to hamstring ACL reconstruction in young active patients significantly decreased the rate of graft failure.

Focusing more closely on secondary meniscal injuries, we found a rate of 1.9% in the 4HT + ALL group versus 9.9% in the 4HT group. Secondary meniscal lesions after ACL reconstruction have been investigated thoroughly. Sonnerly-Cottet et al. [49] found that the probability of failure

of medial meniscal repairs was > 2 times lower patients undergoing ACL + ALL reconstruction compared to ACL reconstruction alone (HR = 0.443 [95% CI: 0.218–0.866]). Two systematic reviews of 13 and 21 studies [33, 56] found a reoperation rate for secondary meniscal lesions of 26.9% and 14.2%, respectively, 5 years after ACL reconstruction, which is higher than the two groups of this study. Another study by Balendra et al. [5] found a rate of secondary meniscal injury of 16.4% in professional football players, 2 years after ACL reconstruction.

Table 5 Return to sport, subjective outcomes and PROMs among the whole cohort and the two surgical groups at final follow-up

Outcome	Total population (N=203)	HT (N=101)	HT + ALL (N=102)	p-value
Return to same sport	140 (75.3)	68 (75.6)	72 (75)	n.s.
Return to same sport at the same level	88 (47.3)	38 (42.2)	50 (52)	n.s.
Occasional pain	68 (38.6)	34 (37.8)	34 (35.4)	n.s.
Stiffness	41 (22)	20 (22.2)	21 (21.9)	n.s.
Instability	33 (17.7)	15 (16.7)	18 (18.7)	n.s.
Hardware discomfort	14 (7.5)	8 (8.9)	6 (6.3)	n.s.
Swelling	24 (13)	11 (12.3)	13 (13.6)	n.s.
IKDC score				n.s.
Mean \pm SD	82.7 \pm 14.3	83.3 \pm 14.3	82 \pm 14.4	
Range (min–max)	(37–100)	(37–100)	(42–100)	
ACL-RSI score				n.s.
Mean \pm SD	68.6 \pm 22.9	69.8 \pm 23.5	67.4 \pm 22.4	
Range (min–max)	(8–100)	(8–100)	(16–100)	
Tegner activity scale				n.s.
Mean	7	7	7	
Range (min–max)	(3–10)	(3–10)	(4–10)	
Lysholm knee score				n.s.
Mean (\pm SD)	85.8 \pm 15.9	86.4 \pm 15.2	86 \pm 16.8	
Range (min–max)	(44–100)	(44–100)	(46–100)	

Values shown are *n* (%) unless stated otherwise

HT, hamstring tendon; HT + ALL, hamstring tendon + anterolateral ligament; IKDC, International Knee Documentation Committee score; ACL-RSI, anterior cruciate ligament-return to sport after injury scale; SD, standard deviation

p value was not significant for any of the outcomes ($p > 0.05$)

Regarding the postoperative outcomes and PROMs, these were not significantly different between the two surgical procedures and this is consistent with the results of Sonnery-Cottet et al. [50] who found no difference in IKDC, Lysholm knee score and Tegner activity score between three grafts: 4HT, HT + ALL and B-PT-B.

No difference in return to sport was found between the two groups (42.2% of patients in the 4HT group returned to the same sport at the same level versus 52% of patients in the 4HT + ALL group). Webster et al. [55] reported that 88% of patients < 20-years-old returned to high-risk sports 60 months after surgery, while Balendra et al. [5] reported that 96.1% of professional footballers returned to play and 90.1% returned to the same level of play. In a systematic review of ten studies including patients < 15-years-old, Morvan et al. [32] found that a return to sport was possible for 91.7% of individuals, and in four of these studies, 61–89% of patients returned to the same level and 42% returned to a competitive level, after a mean follow-up of 7.9 years. The rate of patients returning to sport in the current study cohort is lower than that in the literature, but according to Kyritsis et al. and Webster et al. [25, 54], we authorised a return to sport 6 months after surgery only if there was 100% muscular recovery, no difference in the jump test, an

ACL-RSI score > 85% and side-to-side laxity < 5 mm. It is possible that the strict criteria for a return to sport used in this study were responsible for the low rate of return to sport. In addition, Coquard et al. [13] observed that adding an ALL reconstruction to a HT ACL graft does not delay functional recovery and return to sport.

Some surgeons are concerned about doing ALL reconstructions because of a presumed higher risk of complications. In the current study, there was no difference in subjective outcomes such as occasional knee pain, hardware problems, stiffness and swelling 2 years after surgery. Furthermore, there was also no difference regarding cyclops syndrome.

Although the operating time was significantly longer in the 4HT + ALL group, it did not lead to more infections; there were only two infections, both in the 4HT group, which occurred prior to 2017 and before the use of the vancomycin-soaked compresses [22, 35]. However, there were three cases of hardware removal in the 4HT + ALL group, all related to ALL fixation, versus none in the 4HT group. According to previous experience, these hardware discomforts originate essentially from the femoral-ALL fixation screw, which must be well buried in the bone in order not to rub against the iliotibial band. These results

are, therefore, reassuring when it comes to the supposed risks of anatomic ALL reconstruction and are in accordance with the results of Sonnery-Cottet et al. [48].

The ideal indication for 4HT + ALL has yet to be established, namely which patients could benefit from it and which can do without it? In our opinion, the indication for ALL reconstruction is specific to each patient and should be based on clinical examination and laxity findings. According to Cavaignac et al. [9], it is possible to screen an ALL lesion by ultrasound in the operating room before making the first incision. This may be a way forward to refine the indications and would also allow a more reproducible placement of the femoral tunnel according to Castoldi et al. [8].

This study has several limitations. First, it was retrospective in design and the lack of randomisation prevents us from ruling out selection bias. Follow-up was only 3 years, but the majority of graft failures occur during the first 2 years after surgery and the mean time to failure of 1.9 years is consistent with the literature [57]. Follow-up was also longer in the 4HT group, which may account for some of the differences in failure rates, but the mean time to failure was not different between the two groups. Further multicentre, randomised, controlled studies are needed to refine the indications for anterolateral extraarticular procedures and to confirm the current results, which demonstrate better outcomes with combined ACL + ALL reconstruction in this high-risk population.

Conclusion

In young athletes, ALL reconstruction reduces the graft failure rate by half and significantly reduces the incidence of secondary meniscal lesions. The two independent risk factors for reoperation due to graft failure or secondary meniscal lesions were the absence of an ALL graft and preoperative side-to-side laxity > 3 mm. Combined ACL and ALL reconstruction should be preferred in young patients who participate in pivoting sports.

Acknowledgements None.

Author contributions PL: design, data acquisition, statistical analysis, interpretation of data, writing and revision of the manuscript. AD: data acquisition, interpretation of data, writing and revision of the manuscript. NB: data acquisition, writing and revision of the manuscript. AB: interpretation of data, supervision, writing and revision of the manuscript. NG: design, data acquisition, statistical analysis, interpretation of data, writing and revision of the manuscript. All authors have reviewed and approve the written version of the manuscript.

Funding The authors did not receive support from any organization for the submitted work.

Declarations

Conflict of interest The authors have no conflicts of interest to declare that are relevant to the content of this article.

Ethical approval This retrospective study involving human participants was in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. The study was approved by the local Institutional Review Board (No. 12-2019-12).

Informed consent All patients gave informed consent to participate to the study.

References

1. Alkhalaf FNA, Hanna S, Alkhalidi MSH, Alenezi F, Khaja A (2021) Autograft diameter in ACL reconstruction: size does matter. *SICOT-J* 7:16
2. Andernord D, Desai N, Björnsson H, Ylander M, Karlsson J, Samuelsson K (2015) Patient predictors of early revision surgery after anterior cruciate ligament reconstruction: a cohort study of 16,930 patients with 2-year follow-up. *Am J Sports Med* 43:121–127
3. Ariel de Lima D, Helito CP, Lacerda de Lima L, de Castro SD, Costa Cavalcante ML, Dias Leite JA (2019) Anatomy of the anterolateral ligament of the knee: a systematic review. *Arthroscopy* 35:670–681
4. Ariel de Lima D, Helito CP, Lacerda de Lima L, Dias Leite JA, Costa Cavalcante ML (2019) Study of the nerve endings and mechanoreceptors of the anterolateral ligament of the knee. *Arthroscopy* 35:2918–2927
5. Balendra G, Jones M, Borque KA, Willinger L, Pinheiro VH, Williams A (2021) Factors affecting return to play and graft rupture after primary ACL reconstruction in professional footballers. *Knee Surg Sports Traumatol Arthrosc*. <https://doi.org/10.1007/s00167-021-06765-8>
6. Bourke HE, Salmon LJ, Waller A, Patterson V, Pinczewski LA (2012) Survival of the anterior cruciate ligament graft and the contralateral ACL at a minimum of 15 years. *Am J Sports Med* 40:1985–1992
7. Boutsidiadis A, Brossard P, Panisset J-C, Graveleau N, Barth J (2017) Minimally invasive combined anterior and anterolateral stabilization of the knee using hamstring tendons and adjustable-loop suspensory fixation device: surgical technique. *Arthrosc Tech* 6:e419–e425
8. Castoldi M, Cavaignac M, Marot V, Reina N, Mouarbes D, Berard E, Cavaignac E (2022) Femoral positioning of the anterolateral ligament graft with and without ultrasound location of the lateral epicondyle. *Am J Sports Med* 50:415–422
9. Cavaignac E, Laumond G, Reina N, Wytrykowski K, Murgier J, Faruch M, Chiron P (2018) How to test the anterolateral ligament with ultrasound. *Arthrosc Tech* 7:e29–e31
10. Claes S, Hermie L, Verdonk R, Bellemans J, Verdonk P (2013) Is osteoarthritis an inevitable consequence of anterior cruciate ligament reconstruction? A meta-analysis. *Knee Surg Sports Traumatol Arthrosc* 21:1967–1976
11. Claes S, Vereecke E, Maes M, Victor J, Verdonk P, Bellemans J (2013) Anatomy of the anterolateral ligament of the knee. *J Anat* 223:321–328
12. Cooper JD, Wang W, Prentice HA, Funahashi TT, Maletis GB (2019) The Association between tibial slope and revision anterior cruciate ligament reconstruction in patients ≤21 years old: a

- matched case-control study including 317 revisions. *Am J Sports Med* 47:3330–3338
13. Coquard M, Carrozzo A, Saithna A, Vigne G, Le Guen M, Fournier Y, Hager J-P, Vieira TD, Sonnery-Cottet B (2022) Anterolateral ligament reconstruction does not delay functional recovery, rehabilitation, and return to sport after anterior cruciate ligament reconstruction: a matched-pair analysis from the SANTI (Scientific ACL Network International) Study Group. *Arthrosc Sports Med Rehabil* 4:e9–e16
 14. Cruz AI, Beck JJ, Ellington MD, Mayer SW, Pennock AT, Stinson ZS, VandenBerg CD, Barrow B, Gao B, Ellis HB (2020) Failure rates of autograft and allograft ACL reconstruction in patients 19 years of age and younger: a systematic review and meta-analysis. *JB JS Open Access* 5(e20):00106
 15. Ferretti A, Monaco E, Redler A, Argento G, De Carli A, Saithna A, Helito PVP, Helito CP (2019) High prevalence of anterolateral ligament abnormalities on MRI in knees with acute anterior cruciate ligament injuries: a case-control series from the SANTI study group. *Orthop J Sports Med* 7:2325967119852916
 16. Firth AD, Bryant DM, Litchfield R, McCormack RG, Heard M, MacDonald PB, Spalding T, Verdonk PCM, Peterson D, Bardana D, Rezansoff A, STABILITY Study Group, Getgood AMJ, Willits K, Birmingham T, Hewison C, Wanlin S, Pinto R, Martindale A, O'Neill L, Jennings M, Daniluk M, Boyer D, Zomar M, Moon K, Moon R, Fan B, Mohan B, Buchko GM, Hiemstra LA, Kerslake S, Tynedal J, Stranges G, Mcrae S, Gullett L, Brown H, Legary A, Longo A, Christian M, Ferguson C, Mohtadi N, Barber R, Chan D, Campbell C, Garven A, Pulsifer K, Mayer M, Simunovic N, Duong A, Robinson D, Levy D, Skelly M, Shanmugaraj A, Howells F, Tough M, Thompson P, Metcalfe A, Asplin L, Dube A, Clarkson L, Brown J, Bolsover A, Bradshaw C, Belgrove L, Milan F, Turner S, Verdugo S, Lowe J, Dunne D, McGowan K, Suddens C-M, Declerq G, Vuylsteke K, Van Haver M (2022) Predictors of Graft failure in young active patients undergoing hamstring autograft anterior cruciate ligament reconstruction with or without a lateral extra-articular tenodesis: the stability experience. *Am J Sports Med* 50:384–395
 17. Frank JS, Gambacorta PL (2013) Anterior cruciate ligament injuries in the skeletally immature athlete: diagnosis and management. *J Am Acad Orthop Surg* 21:78–87
 18. Geffroy L, Lefevre N, Thevenin-Lemoine C, Peyronnet A, Lakhil W, Fayard JM, Chotel F, French Arthroscopy Society (2018) Return to sport and re-tears after anterior cruciate ligament reconstruction in children and adolescents. *Orthop Traumatol Surg Res* OTSR 104:S183–S188
 19. Hewison CE, Tran MN, Kaniki N, Remtulla A, Bryant D, Getgood AM (2015) Lateral extra-articular tenodesis reduces rotational laxity when combined with anterior cruciate ligament reconstruction: a systematic review of the literature. *Arthroscopy* 31:2022–2034
 20. Hoshino Y, Fu FH, Irrgang JJ, Tashman S (2013) Can joint contact dynamics be restored by anterior cruciate ligament reconstruction? *Clin Orthop Relat Res* 471:2924–2931
 21. Hyun Ahn J, Jun Koh I, McGarry MH, Patel NA, Lin CC, Lee TQ (2022) Synergistic effect of the anterolateral ligament and capsule injuries on the knee laxity in anterior cruciate ligament injured knees: a cadaveric study. *Orthop Traumatol Surg Res* OTSR 2022:103224
 22. Jefferies JG, Aithie JMS, Spencer SJ (2019) Vancomycin-soaked wrapping of harvested hamstring tendons during anterior cruciate ligament reconstruction. A review of the “vancomycin wrap.” *Knee* 26:524–529
 23. Kamath GV, Murphy T, Creighton RA, Viradia N, Taft TN, Spang JT (2014) Anterior cruciate ligament injury, return to play, and reinjury in the elite collegiate athlete: analysis of an NCAA division I cohort. *Am J Sports Med* 42:1638–1643
 24. Kamien PM, Hydrick JM, Replogle WH, Go LT, Barrett GR (2013) Age, graft size, and Tegner activity level as predictors of failure in anterior cruciate ligament reconstruction with hamstring autograft. *Am J Sports Med* 41:1808–1812
 25. Kyritsis P, Bahr R, Landreau P, Miladi R, Witvrouw E (2016) Infographic. Avoid ACL graft rupture. Meet discharge criteria. *Br J Sports Med* 50:952
 26. LaPrade RF (2016) Editorial commentary: it is all about how one defines the anatomy. *Arthroscopy* 32:849–850
 27. Lee DW, Kim JG, Cho SI, Kim DH (2019) Clinical outcomes of isolated revision anterior cruciate ligament reconstruction or in combination with anatomic anterolateral ligament reconstruction. *Am J Sports Med* 47:324–333
 28. Lee DW, Lee JK, Kwon SH, Moon SG, Cho SI, Chung SH, Kim JG (2021) Adolescents show a lower healing rate of anterolateral ligament injury and a higher rotational laxity than adults after anterior cruciate ligament reconstruction. *Knee* 30:113–124
 29. Lefevre N, Bohu Y, Naouri JF, Klouche S, Herman S (2014) Validity of GNRB® arthrometer compared to Telos™ in the assessment of partial anterior cruciate ligament tears. *Knee Surg Sports Traumatol Arthrosc* 22:285–290
 30. Leys T, Salmon L, Waller A, Linklater J, Pinczewski L (2012) Clinical results and risk factors for reinjury 15 years after anterior cruciate ligament reconstruction: a prospective study of hamstring and patellar tendon grafts. *Am J Sports Med* 40:595–605
 31. Morgan MD, Salmon LJ, Waller A, Roe JP, Pinczewski LA (2016) Fifteen-year survival of endoscopic anterior cruciate ligament reconstruction in patients aged 18 years and younger. *Am J Sports Med* 44:384–392
 32. Morvan A, Bouguennec N, Graveleau N (2020) ACL injuries before 15 years of age: could the young become an athlete? *Arch Orthop Trauma Surg*. <https://doi.org/10.1007/s00402-020-03404-8>
 33. Nepple JJ, Dunn WR, Wright RW (2012) Meniscal repair outcomes at greater than five years: a systematic literature review and meta-analysis. *J Bone Joint Surg Am* 94:2222–2227
 34. Nwachukwu BU, Voleti PB, Berkanish P, Chang B, Cohn MR, Williams RJ, Allen AA (2017) Return to play and patient satisfaction after ACL reconstruction: study with minimum 2-year follow-up. *J Bone Joint Surg Am* 99:720–725
 35. Offerhaus C, Balke M, Hente J, Gehling M, Blendl S, Höher J (2019) Vancomycin pre-soaking of the graft reduces postoperative infection rate without increasing risk of graft failure and arthrofibrosis in ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 27:3014–3021
 36. Perelli S, Morales-Avalos R, Formagnana M, Rojas-Castillo G, Serranocolí G, Monllau JC (2022) Lateral extraarticular tenodesis improves stability in non-anatomic ACL reconstructed knees: in vivo kinematic analysis. *Knee Surg Sports Traumatol*. <https://doi.org/10.1007/s00167-021-06854-8>
 37. Persson A, Fjeldsgaard K, Gjertsen J-E, Kjellsen AB, Engebretsen L, Hole RM, Fevang JM (2014) Increased risk of revision with hamstring tendon grafts compared with patellar tendon grafts after anterior cruciate ligament reconstruction: a study of 12,643 patients from the Norwegian Cruciate Ligament Registry, 2004–2012. *Am J Sports Med* 42:285–291
 38. Rayes J, Ouanezar H, Haidar IM, Ngbilo C, Fradin T, Vieira TD, Freychet B, Sonnery-Cottet B (2022) Revision anterior cruciate ligament reconstruction using bone-patellar tendon-bone graft combined with modified lemaire technique versus hamstring graft combined with anterolateral ligament reconstruction: a clinical comparative matched study with a mean follow-up of 5 years from the SANTI study group. *Am J Sports Med* 50:395–403
 39. Rhatomy S, Ariyanto MW, Fiolin J, Dilogo IH (2022) Comparison of clinical outcomes between isolated ACL reconstruction and combined ACL with anterolateral ligament reconstruction: a

- systematic review and meta-analysis. *Eur J Orthop Surg Traumatol*. <https://doi.org/10.1007/s00590-021-03194-8>
40. Robert H, Nouveau S, Gageot S, Gagnière B (2009) A new knee arthrometer, the GNRB: experience in ACL complete and partial tears. *Orthop Traumatol Surg Res OTSR* 95:171–176
 41. Rochongar G, Cucurulo T, Ameline T, Potel JF, Dalmay F, Pujol N, Sallé de Chou É, Lutz C, Ehkirch FP, Le Henaff G, Laporte C, Seil R, Gunepin F-X, Sonnery-Cottet B (2015) Meniscal survival rate after anterior cruciate ligament reconstruction. *Orthop Traumatol Surg Res* 101:S323–S326
 42. Roessler PP, Schüttler KF, Heyse TJ, Wirtz DC, Efe T (2016) The anterolateral ligament (ALL) and its role in rotational extra-articular stability of the knee joint: a review of anatomy and surgical concepts. *Arch Orthop Trauma Surg* 136:305–313
 43. Rugg CM, Pitcher AA, Allen C, Pandya NK (2020) Revision ACL reconstruction in adolescent patients. *Orthop J Sports Med* 8:2325967120953337
 44. Song G-Y, Hong L, Zhang H, Zhang J, Li Y, Feng H (2016) Clinical outcomes of combined lateral extra-articular tenodesis and intra-articular anterior cruciate ligament reconstruction in addressing high-grade pivot-shift phenomenon. *Arthroscopy* 32:898–905
 45. Sonnery-Cottet B, Daggett M, Fayard J-M, Ferretti A, Helito CP, Lind M, Monaco E, de Pádua VBC, Thauinat M, Wilson A, Zaffagnini S, Zijl J, Claes S (2017) Anterolateral ligament expert group consensus paper on the management of internal rotation and instability of the anterior cruciate ligament—deficient knee. *J Orthop* 18:91–106
 46. Sonnery-Cottet B, Freychet B, Murphy CG, Pupim BHB, Thauinat M (2014) Anterior cruciate ligament reconstruction and preservation: the single-antemedial bundle biological augmentation (SAMBBA) technique. *Arthrosc Tech* 3:e689–e693
 47. Sonnery-Cottet B, Lutz C, Daggett M, Dalmay F, Freychet B, Niglis L, Imbert P (2016) The involvement of the anterolateral ligament in rotational control of the knee. *Am J Sports Med* 44:1209–1214
 48. Sonnery-Cottet B, Pioger C, Vieira TD, Franck F, Kajetanek C, Fayard J-M, Thauinat M, Saithna A (2020) Combined ACL and anterolateral reconstruction is not associated with a higher risk of adverse outcomes: preliminary results from the SANTI randomized controlled trial. *Orthop J Sports Med* 8:2325967120918490
 49. Sonnery-Cottet B, Saithna A, Blakeney WG, Ouanezar H, Borade A, Daggett M, Thauinat M, Fayard J-M, Delaloye J-R (2018) Anterolateral ligament reconstruction protects the repaired medial meniscus: a comparative study of 383 anterior cruciate ligament reconstructions from the SANTI study group with a minimum follow-up of 2 years. *Am J Sports Med* 46:1819–1826
 50. Sonnery-Cottet B, Saithna A, Cavalier M, Kajetanek C, Temponi EF, Daggett M, Helito CP, Thauinat M (2017) Anterolateral ligament reconstruction is associated with significantly reduced ACL graft rupture rates at a minimum follow-up of 2 years: a prospective comparative study of 502 patients from the SANTI study group. *Am J Sports Med* 45:1547–1557
 51. Sonnery-Cottet B, Saithna A, Quelard B, Daggett M, Borade A, Ouanezar H, Thauinat M, Blakeney WG (2019) Arthrogenic muscle inhibition after ACL reconstruction: a scoping review of the efficacy of interventions. *Br J Sports Med* 53:289–298
 52. Webster KE, Feller JA (2016) Exploring the high reinjury rate in younger patients undergoing anterior cruciate ligament reconstruction. *Am J Sports Med* 44:2827–2832
 53. Webster KE, Feller JA, Kimp AJ, Whitehead TS (2018) Revision anterior cruciate ligament reconstruction outcomes in younger patients: medial meniscal pathology and high rates of return to sport are associated with third ACL injuries. *Am J Sports Med* 46:1137–1142
 54. Webster KE, Feller JA, Lambros C (2008) Development and preliminary validation of a scale to measure the psychological impact of returning to sport following anterior cruciate ligament reconstruction surgery. *Phys Ther Sport* 9:9–15
 55. Webster KE, Feller JA, Leigh WB, Richmond AK (2014) Younger patients are at increased risk for graft rupture and contralateral injury after anterior cruciate ligament reconstruction. *Am J Sports Med* 42:641–647
 56. Westermann RW, Duchman KR, Amendola A, Glass N, Wolf BR (2017) All-Inside Versus Inside-Out Meniscal Repair With Concurrent Anterior Cruciate Ligament Reconstruction: A Meta-regression Analysis. *Am J Sports Med* 45:719–724
 57. Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD (2016) Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Am J Sports Med* 44:1861–1876
 58. Yamazaki J, Muneta T, Ju Y-J, Koga H, Morito T, Sekiya I (2013) The kinematic analysis of female subjects after double-bundle anterior cruciate ligament reconstruction during single-leg squatting. *J Orthop* 18:284–289
 59. Yoo JH, Chang CB, Shin KS, Seong SC, Kim TK (2008) Anatomical references to assess the posterior tibial slope in total knee arthroplasty: a comparison of 5 anatomical axes. *J Arthroplasty* 23:586–592

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.